

1 **The role of extra-striate areas in conscious motor behavior: a**
2 **registered report with Fast-Optical Imaging**

3 Elisabetta Colombari^a, Giorgia Parisi^a, Sonia Mele^a, Chiara Mazzi^a and Silvia
4 Savazzi^a

5
6 ^a Perception and Awareness (PandA) Laboratory, Department of Neuroscience, Biomedicine and
7 Movement Sciences, University of Verona, Strada le Grazie 8, Verona, Italy

8
9

10 **Abstract**

11 Disclosing the brain areas responsible for the emergence of visual awareness and their timing of
12 activation represents one of the major challenges in consciousness research. In particular, isolating
13 the neural processes strictly related to consciousness from concurrent neural dynamics either related
14 to prerequisites or post-perceptual processing has long engaged consciousness research. In this
15 framework, the present study aims at unravelling the spatio-temporal dynamics underlying conscious
16 vision by adopting a ~~peculiar~~ **distinctive** experimental design in which both awareness and motor
17 response are manipulated, allowing the segregation of neural activity strictly related to awareness
18 from response-related mechanisms. To this aim, we will employ a GO/NOGO detection task, in which
19 participants will respond or withhold responding according to the experimental condition. Critically,
20 during the performance of the task, participants' brain activity will be recorded by means of Event-
21 Related Optical Signal (EROS) technique, which provides accurate information about brain functions
22 both from the temporal and spatial point of view, simultaneously. The combination of this
23 experimental design with EROS recording will enable us to pinpoint the neural correlates underlying
24 conscious vision and to disentangle them from processes related to the response. In addition, by
25 coupling conventional EROS analysis with Granger Causality analysis, we will be able to clarify the
26 potential interplay between consciousness-related extra-striate areas and response-related motor
27 areas.

28 1. Introduction

29 Consciousness, namely the set of subjective experiences we have when we are awake, is one of the
30 most intriguing topics debated in neuroscience research. In particular, the search for its neural
31 correlates (NCC) has permeated the literature in recent decades. In broad strokes, one of the most
32 widely used approaches to assess such NCCs involves contrasting brain activity occurring when a
33 visual stimulus enters consciousness with brain activity occurring when the same stimulus does not
34 reach awareness. This renowned paradigm is known as *contrastive analysis* (Aru *et al.*, 2012) and
35 has been frequently combined with electrophysiological recording or functional neuroimaging,
36 leading to numerous and dissimilar results (Förster *et al.*, 2020). Indeed, the interpretations of spatio-
37 temporal dynamics underlying conscious vision are among the most disparate. ERP studies propose
38 two possible electrophysiological markers as correlates of visual awareness: an earlier occipito-
39 temporal negative deflection (i.e., Visual Awareness Negativity – VAN) detectable 200 ms after the
40 presentation of the stimulus, and a later positivity (i.e., Late Positivity – LP) widespread over centro-
41 parietal regions, peaking 300-500 ms after the stimulus onset (Koivisto & Revonsuo, 2010). However,
42 ~~the electrophysiological signature/s characterizing which one represents the true signature of~~
43 ~~conscious vision is~~ **has still to be elucidated** ~~under debate~~. This may be attributed to one of the main
44 limitations of the contrastive analysis, which is represented by its ineffectiveness in dissociating the
45 true NCC (i.e., the set of neural correlates necessary and sufficient to enable consciousness) from
46 concurrent neural dynamics either related to prerequisites or post-perceptual processing (Aru *et al.*,
47 2012). In most prior studies aiming at identifying such NCCs, participants were asked to make
48 judgments about their experience. However, such an operation could lead to confounding neural
49 processes related to the task, not strictly to awareness per se.

50 For this reason, in an effort to disentangle the proper correlates of consciousness from neural activity
51 related to the response, no-report paradigms have been employed. In this framework, no-report
52 paradigms, where participants are not requested to perform any tasks or to provide any judgments
53 about their perceptual experience, represent an advantageous tool to dissociate the neural processes
54 strictly related to consciousness from subsequent processes related to the required response (Tsuchiya
55 *et al.*, 2015; Hatamimajoumerd *et al.*, 2022). Studies employing this kind of **paradigm with different**
56 **techniques such as EEG and fMRI** concluded that LP is highly modulated by several different
57 cognitive processes occurring at later stages of processing (Mazzi *et al.*, 2020; Schlossmacher *et al.*,
58 2020; Dembski *et al.*, 2021; Kronemer *et al.*, 2022), **as well as by the task relevance of the stimulus**
59 **(Makeig & Jung, 2000; Pitts *et al.*, 2014; Shafto & Pitts, 2015; Schelonka *et al.*, 2017; Dellert *et al.*,**
60 **2021; Hense *et al.*, 2024). By contrast, the role of response requirements, as well as that of attention,**
61 **on the VAN are still debated** ~~seems not to be sensitive to the task or the response (Cohen *et al.*,~~

62 ~~2020~~) as different studies have reported both positive (e.g., Bola & Doradzińska, 2021; Dellert *et al.*,
63 2021; Doradzińska & Bola, 2024) and negative (e.g., Koivisto *et al.*, 2006; Cohen *et al.*, 2020; Dellert
64 *et al.*, 2022; Ciupińska *et al.*, 2024) results. ~~Indeed,~~ Interestingly, in a study published in 2016 by
65 Koivisto and colleagues (Koivisto *et al.*, 2016), authors successfully dissociated ERP correlates of
66 visual awareness from those related to ~~post-perceptual mechanisms~~ response, disclosing that VAN
67 was not modulated by response requirements. The authors adopted a particular partial-report
68 paradigm in which participants were sometimes asked to provide a report by pressing a response
69 button when they were aware of the stimulus and sometimes to withhold responding in case of
70 awareness. They found that, while the amplitude of LP was modulated by the response (i.e., it was
71 greater in trials where participants were asked to respond in case of awareness, compared to the Aware
72 condition where they were asked to withhold responding), VAN did not change depending on task
73 requirements. This allowed Koivisto and colleagues to advocate for an early onset of visual
74 awareness: the phenomenal content of a visual experience, indeed, takes place before LP, more
75 specifically in the temporal window of VAN.

76 Several pieces of evidence are consistent in considering VAN as the electrophysiological signature
77 of phenomenal consciousness (Koivisto *et al.*, 2008; Railo *et al.*, 2015), ~~while~~ the localization of its
78 neural generator still remains open. In this regard, previous MEG source localization studies (Vanni
79 *et al.*, 1996; Liu *et al.*, 2012) identified the Lateral Occipital Complex (LOC), an extra-striate visual
80 areas traditionally associated with objects recognition, as the generator of VAN. The same result was
81 achieved in a recent work which aimed at unravelling the spatio-temporal dynamics underlying
82 conscious vision (Colombari *et al.*, 2024). In such study, participants were asked to perform a
83 discrimination task on the orientation of a tilted Gabor patch while their brain activity was recorded
84 first with EEG and then with Fast Optical Imaging. This allowed authors to identify the exact temporal
85 window of VAN and LP and then, by taking advantage of the peculiarity of Fast Optical Imaging of
86 achieving both temporal and spatial accurate information (Gratton & Corballis, 1995; Gratton &
87 Fabiani, 2010; Baniqued *et al.*, 2013), to investigate the spatio-temporal unfolding of brain activity
88 occurring in these predetermined time windows. Authors contrasted activity of Aware trials (i.e.,
89 trials in which participants reported to perceive the orientation of the stimulus) with activity of
90 Unaware ones and observed a sustained activation of LOC in the VAN temporal window, consistently
91 with the above-mentioned MEG studies. More interestingly, they observed that, only when the
92 stimulus crossed the threshold of consciousness, activity in extra-striate visual areas triggered
93 subsequent activation of motor areas, although motor response was required in both Aware and
94 Unaware conditions. Authors tried to interpret this unexpected finding by ascribing it to the selection
95 of the correct response, that could be provided in the Aware trials only where participants consciously

96 perceived the stimulus. Indeed, in Aware trials participants had to press a specific button on the
97 response box (to provide the correct answer about the orientation of the Gabor patch), while when
98 the stimulus was unseen (i.e., Unaware trials) they had to respond randomly, by pressing indifferently
99 one of the two response buttons. However, the employed experimental paradigm did not allow the
100 authors to thoroughly investigate this issue. Thus, in order to clarify the interplay between extra-
101 striate areas and motor regions in awareness, in the present study we will adopt a go/no-go detection
102 task (similar to that adopted by Koivisto *et al.*, 2016), while recording participants' brain activity by
103 means of Fast Optical Imaging. Specifically, Event-Related Optical Signal (EROS) technique will be
104 employed. This technique, by shedding near-infrared light through the brain tissues, is able to detect
105 changes in the light scattering properties that are known to be directly related to neural activity, thus
106 providing accurate information about brain functions both from the temporal and spatial point of
107 view, simultaneously (Gratton *et al.*, 1997; Gratton & Fabiani, 1998, 2001). Critically, the study will
108 adopt a peculiar paradigm manipulating both awareness and response. The latter, indeed, will be
109 provided sometimes in the Aware condition (condition Aware-GO/Unaware-NOGO) and sometimes
110 in the Unaware one (condition Aware-NOGO/Unaware-GO). This double manipulation will enable
111 us ~~to~~ to unravel the spatio-temporal unfolding of awareness-related activity, by ~~isolating~~ **disentangling**
112 neural activity ~~strictly~~ related to awareness from response-related mechanisms. **Indeed, in the present**
113 **study, we can investigate the NCCs both when the motor response is required and when no task is**
114 **performed, thus allowing to isolate consciousness effects from the effects related to the task.**
115 **Importantly, the experimental paradigm adopted will enable us ~~and ii)~~ to elucidate the interplay**
116 between extra-striate visual areas and motor areas. Indeed, in addition to conventional EROS
117 analyses, we will perform Granger Causality analysis, in order to disclose the relationship existing
118 among the investigated areas. In broad strokes, Granger analysis allows to move beyond the classical
119 identification of cortical activation provided by EROS analysis by disclosing functional circuits
120 underpinning the investigated brain function (Seth *et al.*, 2015). When coupled with EROS, Granger
121 Causality analysis represents a powerful tool to highlight predictive relationship between activations
122 in the investigated regions of interest (ROI) at different time-points (Parisi *et al.*, 2020).

123 Based on previous literature suggesting that VAN is independent from subjective report (Koivisto *et*
124 *al.*, 2016; Ye *et al.*, 2024) and LOC represents the cortical generator or VAN (Liu *et al.*, 2012;
125 Colombari *et al.*, 2024), we expect Aware trials to elicit early greater activation of LOC,
126 independently of the response requirement. Moreover, by combining EROS conventional analysis
127 with Granger Causality analysis, and manipulating both awareness and motor response, we aim to
128 highlight potential interplay between consciousness-related extra-striate areas and response-related
129 motor areas both when the motor response is required and when it has to be inhibited.

130 2. Methods

131 2.1 Ethics Information

132 The study is approved by the local Ethics Committee (Prog.171CESC) and it will be conducted in
133 accordance with the principles laid down in the 2013 Declaration of Helsinki ~~and~~. Participants will
134 be recruited from the University of Verona community, by means of printed flyers displayed on notice
135 boards at different University of Verona sites and through advertisements on social media. Each
136 participant will be fully informed about the modalities of the study before taking part in the
137 experiment and written informed consent will be signed. In addition, participants will receive
138 compensation for their participation and will be debriefed after the conclusion of the experiment.

139 2.2 Participants

140 We will recruit healthy adults, right-handed (as assessed by means of the standard handedness
141 inventory *Edinburgh Handedness Questionnaire*; Oldfield, 1971) and aged between 18 and 50 years
142 old. All of them will have to report normal or corrected-to-normal vision, no history of neurological
143 or psychiatric disorders and no contraindications to MRI. The study will be conducted at the Panda
144 lab of the University of Verona (Italy).

145 2.2.1 Sample size estimation

146 Since in EROS literature no previous studies report the effect size because of technical constraints of
147 the employed dedicated software, an a priori statistical sample size estimation for the present study
148 is not achievable based on EROS data. For this reason, we first based our sample size estimation on
149 a review of the existing EROS literature (see Supplementary Table 1 at
150 https://osf.io/ebfu3/?view_only=9ec2e6bf32ba4a8bb8b858639ec40a59) (GRATTON *et al.*, 1995;
151 Gratton *et al.*, 1997, 2000, 2001; Gratton & Fabiani, 2003; Wolf *et al.*, 2003; Low *et al.*, 2006; Tse
152 & Penney, 2007; Medvedev *et al.*, 2008, 2010; Proulx *et al.*, 2018; Toscano *et al.*, 2018; Parisi *et al.*,
153 2020; Tse *et al.*, 2021; Knight *et al.*, 2024), from which emerges that, on average, EROS studies
154 employ experimental samples composed of ~~42~~ 13 participants (mean 12.944; SD 7.008). Moreover,
155 we decided to estimate the sample size for the present study basing on a previous EEG study
156 employing the same experimental design adopted in the present study (Koivisto *et al.*, 2016). The
157 estimated sample size for research questions Q1 (i.e., “Can we replicate Colombari *et al.*, 2024
158 findings showing that LOC is an NCC?”) and Q2 (i.e., “Is the activity in LOC independent from the
159 response?”) was calculated with G-Power software (v. 3.1.9.7), with a power of 90% and a level of
160 significance of 2%. To estimate the power needed to detect the effect of awareness (aware vs. unaware
161 trials), we considered the significant main effect of awareness of a within subjects repeated measures

162 ANOVA ($F(1,14) = 17.06$, $P = 0.001$, $\eta_p^2 = 0.55$) carried out in Koivisto et al.(2016). The estimated
163 sample size resulted in 15 participants (critical $F = 6.887$; actual power = 0.918). Since EROS signal-
164 to-noise ratio is lower than that of EEG, we will increase our final sample to 26 24 participants.
165 Considering that the estimated sample size for this study ($n = 26 24$) is the double of the typical sample
166 size of EROS studies present in literature, the same estimated sample size seems to be also adequate
167 to answer research questions Q3 (i.e., “Does consciousness modulate activation of motor areas in a
168 detection task?”) and Q4 (i.e., “Does consciousness modulate activation of motor areas in ABSENCE
169 of motor response?”).

170 2.2.2 Exclusion Criteria

171 As better specified in section 2.3, before getting involved in the study, participants will undergo a
172 perceptual threshold assessment, in order to identify the proper stimulus to be employed in the main
173 experiment. To be enrolled in the study, participants will have to successfully complete this session.
174 The criterion used is that one of the stimuli presented during the threshold assessment will have to
175 be acknowledged as perceived a minimum of 25%, a maximum of 75%, or closest to the 50% of the
176 times the 50% of the times (i.e., at perceptual threshold level). If no stimulus results at the threshold
177 level, the participant should not be enrolled in the study.

178 In addition, participants who will not complete all the experimental sessions, as well as participants
179 reporting a level of Awareness superior to 75% or inferior to 25% at the end of the experiment will
180 be excluded from analyses. This is to maintain comparable the number of trials in the two
181 experimental conditions (i.e., Aware and Unaware) and to ensure a reliable EROS activity (because
182 of its relatively low signal-to-noise ratio, EROS needs a high number of trials per condition, in
183 order to compute statistics). Moreover, participants whose behavioral performance will be affected
184 by biases related to the behavioral response (as assessed by catch trial analysis, explained more in
185 detail below) will be excluded from the analyses (see below –Section 2.8.1 Behavioral data for
186 more detailed information on the analysis of catch trials). Finally, participants whose EROS signal
187 could not be detected properly during the experiment (for example because of too dark hair or
188 technical issues) will not also be included in the analyses. In particular, the opacity value (i.e., the
189 product of the scattering and absorption coefficients) will be estimated for each participant. Based
190 on this value, it is possible to judge the quality of the signal for each participant, independently
191 from the experimental condition. Opacity values of all participants will be averaged together
192 providing the absorption coefficient to be used when running statistical analysis. Participants whose
193 opacity value is equal to 0 or exceeds three standard deviations of the mean will be excluded from
194 statistical analyses.

195 Importantly, each participant who will be excluded due to the previously mentioned exclusion criteria,
196 will be replaced with the recruitment of another participant. Thus, the number of participants to be
197 recruited will be increased to reach a total of ~~26~~²⁴ analyzed subjects, as specified in section 2.2.1.

198 **2.3 Stimuli**

199 Stimuli will be created by means of a custom-made Matlab script (version R2022b; the MathWorks,
200 Inc., Natick, MA) and resized by means of Photoshop (Adobe Photoshop CC, v2014.0.0). As shown
201 in Figure 1, they will be gray circles (.85 .85 .85 RGB), presented on a black background, with 8 radii
202 equally distanced one from another. One radius (the first one, clockwise) can be slightly thicker than
203 the others (critical trials) or not (catch trials). The thickness of the radius for critical stimuli will be
204 individually assessed for each participant on the basis of a subjective perceptual threshold assessment
205 that will be held before the main experiment.

206 Both in the perceptual threshold assessment and in the main experiment, the stimulus will be
207 presented in the lower right quadrant of the screen, specifically at an eccentricity of 3.5° from the
208 fixation cross along the vertical meridian and of 2° along the horizontal one. This is to allow a left-
209 lateralized EROS montage, as a full-head montage is not achievable in our lab due to technical
210 constraints (i.e., insufficient probes). Moreover, since EROS technique is sensitive to depth, a right-
211 lateralized stimulus ensures that it elicits activity in the left portion of the primary visual cortex, which
212 is known to be anatomically closer to the skull compared to the right one, thus ensuring a better
213 penetration of near-infrared light through brain tissues.

214 **2.4 Perceptual Threshold Assessment**

215 Before starting the experiment, participants will undergo a perceptual threshold assessment, with the
216 aim of identifying, for each participant, the level of thickness of the critical radius so that it results to
217 be perceived as thicker 50% of the times. To this aim, stimuli with different levels of radius thickness
218 will be randomly presented and the subjective perceptual threshold will be measured using the method
219 of constant stimuli. Specifically, 9 levels of radius thickness will be presented. The range of stimuli
220 to be used in the perceptual threshold assessment will be selected based on the results of a pilot
221 experiment in which participants were asked to perform the same task employed in the perceptual
222 threshold assessment while presented with a wider range of radius thickness. This will allow us to
223 identify a smaller range of optimal stimuli to be presented thus excluding a range of stimuli whose
224 thickness was almost never or always reported by participants. Each level of radius thickness will be
225 presented 5 times per block, for a total of 8 blocks. Thus, all the stimuli, as well as the catch stimulus,
226 will be presented 40 times each. Participants will be asked to press the spacebar as soon as they detect

227 the stimulus with a thicker radius. The stimulus identified as perceived a minimum of 25%, a
228 maximum of 75%, and closest to 50% of the times 50% of times at the end of the subjective perceptual
229 threshold assessment will be used in the experimental task, together with the catch. The perceptual
230 threshold assessment, as well as the main experiment, will be conducted in a dimly illuminated room
231 and participants will be sitting in front of a 17 in. LCD monitor (resolution 1920x1080, refresh rate
232 of 144 Hz) placed at a viewing distance of 57 cm. Their head will be held in place by means of an
233 adaptable chin rest so that eyes are aligned with the center of the screen. Both the perceptual threshold
234 assessment and the main experiment will be programmed and administered using E-Prime 3.0
235 software (E-Prime Psychology Software Tools Inc., Pittsburgh, PA, USA). Before starting the
236 perceptual threshold assessment, participants will undergo a fixation training (Leung *et al.*, 2009), in
237 order to ensure they will maintain their gaze on the central fixation cross correctly.

238 2.5 Experimental Procedure

239 The experiment will be composed of two identical sessions lasting approximately 3 hours each
240 performed on different days. The first session will be preceded by the assessment of the subjective
241 perceptual threshold, which, in turn, will last around 20 minutes. The two experimental sessions will
242 be identical except for the EROS montages, specifically devised to obtain better coverage of the brain
243 areas of interest. The order of the montages will be counterbalanced across participants, as well as
244 the order of conditions (see below for more detailed information).

245 The task will be a two-conditions go/no-go detection task, similar to that adopted by Koivisto *et al.*,
246 2016, in which participants have to respond in different ways according to the experimental condition
247 (Table 1). In condition “Aware-GO”, they will be asked to press the spacebar on the keyboard as soon
248 as they perceive the thicker radius, and withhold responding when they do not perceive any difference
249 among radii. On the contrary, in condition “Aware-NOGO”, participants will be asked to withhold
250 responding when they perceive a thicker radius, and press the response button when they do not
251 perceive any difference. Each trial will begin with the presentation of a central fixation cross,
252 followed 500 ms later by a sound (1000Hz) presented for 100 ms, notifying participants of the
253 subsequent onset of the stimulus. After a random interval ranging from 500 to 600 ms, the stimulus
254 will be presented for 100 ms in the lower right quadrant of the screen specifically at an eccentricity
255 of 3.5° from the fixation cross along the vertical meridian and of 2° along the horizontal one. After
256 that, participants will be asked to respond according to the experimental condition. Each experimental
257 session will be composed of 24 blocks: 12 blocks for condition Aware-GO/Unaware-NOGO and 12
258 blocks for condition Aware-NOGO/Unaware-GO, counterbalanced across participants according to
259 the order depicted in Table 1. Each block will consist of 50 critical trials and 15 catch trials. The

260 whole experiment will be composed of 48 blocks per participant, for a total of 2400 critical trials and
 261 720 catch trials per participant.

		Awareness	
		yes	no
Response	no	Aware-NOGO	Unaware-NOGO
	yes	Aware-GO	Unaware-GO

262 **Table 1. Experimental conditions.** Both Awareness and Response are manipulated: Awareness is
 263 experimentally manipulated by employing a threshold stimulus, so that sometimes it is consciously perceived
 264 (Aware) and sometimes not (Unaware). Response is manipulated by the task: in condition GO participants are
 265 asked to respond by pressing a key, while in condition NOGO they are asked to withhold responding. The
 266 combination of these two manipulations gives rise to the 4 experimental conditions depicted in the table.

267

Participants	Day 1		Day 2	
	EROS montage 1	Task	EROS montage 2	Task
1	A	GNGG	B	NGGN
2	B	GNGG	A	NGGN
3	A	NGGN	B	GNGG
4	B	NGGN	A	GNGG

268 **Table 2. Counterbalancing of montages and task conditions across participants.** Both EROS montages
 269 and task conditions (G = Aware-GO/Unaware-NOGO; N = Aware-NOGO/Unaware-GO) will be
 270 counterbalanced across participants. In the column “Task”, each letter represents 6 blocks of task. Thus, each
 271 day, participants will perform 12 blocks per condition, for a total of 24 blocks of task per day.

272 2.6 Optical Recording

273 Three synchronized Imagent frequency domain systems (ISS, Inc., Champaign, IL) will be used to
 274 record continuous fast optical data throughout experimental sessions. Each system is equipped with
 275 4 photo-multiplier tubes detectors, for a total of 12 detectors. Near-infrared light (830 nm) will be
 276 delivered from 48 laser diodes on participants’ scalp and it will be modulated at 110 MHz. Each of
 277 12 detectors will receive light from sets of 16 light emitters, multiplexed every 25.6 ms, resulting in
 278 a sampling rate of 39.0625 Hz.

279 To avoid cross-talk between channels, the array of source-detector pairs (i.e., the montage) will be
 280 created by means of a specific program (NOMAD, Near-Infrared Optode Montage Automated
 281 Design) implemented in Matlab, useful to place sources and detectors at optimal distances. In this
 282 experiment, we will set the minimal distance to 17.5 mm and the maximum distance to 50 mm, in

283 order to ensure an extensive coverage of the brain regions of interest both from the spatial and the
284 depth point of view. The distance between the source and the detector of a channel, in fact, determines
285 the depth of the light pathway (Gratton *et al.*, 2000), thus corresponding to the depth of the
286 investigation: namely, longer channels can investigate deeper layers and shorter channels can
287 examine shallower regions.

288 Both light emitters and detectors will be placed on participants head using a custom-built helmet. To
289 minimize interferences, before placing the optical fibers on the head, the hair will be carefully moved
290 with cotton buds, so that the fibers can reach the scalp directly. In order to better adhere to the head
291 of the participant, we will employ two helmets of different sizes: one 55-56 cm large, and one 57-58
292 cm large. For each helmet, we will develop two different montages, so that to provide a dense
293 coverage of the regions of interest (i.e., the left occipital, temporal and parietal cortices, see Figure
294 2). Each montage will consist of the combination of 12 detectors and 48 light emitters, resulting in a
295 total of 192 channels per montage. As mentioned before, each montage will be recorded in a separate
296 session, and the order will be counterbalanced across participants.

297 At the end of each EROS session, the scalp location of each source and detector will be digitized in
298 relation to four fiducial points (i.e., nasion, inion and pre-auricular points) with a neuro-navigation
299 software (SofTaxis, E.M.S., Bologna, Italy) combined with a 3D optical digitizer (Polaris Vicra, NDI,
300 Waterloo, Canada). Afterwards, the digitized scalp locations will be co-registered with each
301 participant's individual MRI, using a dedicated software package (OCP, Optimized Co-registration
302 Package, ~~MATLAB~~ Matlab code) developed by Chiarelli and colleagues (Chiarelli *et al.*, 2015).

303 For this reason, participants will undergo a structural MRI at the Azienda Ospedaliera Universitaria
304 Integrata of Verona (AOUI).

305 **2.7 MRI Acquisition**

306 Participants' individual structural MRI will be acquired by means of a 3 Tesla Philips Ingenia scanner
307 with a 32-channel head RF receive coils. A whole brain high-resolution 3D T1-weighted image (T1w)
308 Turbo-field echo image (1mm-isotropic TE/TR=3.8/8.4 ms, TI=1050 ms) will be acquired.

309 The T1w field of view (240 x 240 x 180 mm) will be large enough to allow for the ears and the entire
310 scalp to be fully included in the image to facilitate later and accurate co-registration with functional
311 data.

312 **2.8 Data Analysis**

313 *2.8.1 Behavioral data*

314 Raw data will be processed by means of custom scripts created on Matlab (the MathWorks, Inc.,
315 Natick, MA). Data will be divided into the 4 experimental conditions (i.e., Aware-GO, Unaware-

316 NOGO, Aware-NOGO, Unaware-GO). For each participant, trials with reaction times lower than 150
317 ms and higher than 3 standard deviations from the mean will be excluded from the analysis. Data will
318 be successively analyzed using Jamovi (version 2.3.28): first, the percentage of Aware and Unaware
319 trials will be calculated, in order to assess that a sufficient amount of trials is present for each
320 condition. Participants presenting more than 75% or less than 25% of Awareness will be discarded
321 from the sample. This is because, in that case, the number of Unaware (or Aware) trials would be
322 insufficient for statistical EROS analysis. EROS technique, indeed, although having a high
323 localization power from both the spatial and temporal point of view, has a relatively low signal-to-
324 noise ratio. For this reason, a high number of trials is required for statistical analysis. Subsequently,
325 reaction times (RTs) will be analyzed for the “GO” conditions, thus paired sample t-tests (two-tailed)
326 will be applied to compare the mean RTs between Aware-GO and Unaware-GO conditions. Finally,
327 to verify that participants are performing the task accurately and that there are no biases related to the
328 response, catch trials will be analyzed. As mentioned above, catch trials are those trials in which all
329 the radii of the stimulus are equally thick, thus no differences in the stimulus are present. In case of
330 catch trials, the participants’ task will be different according to the condition: in the Aware-GO
331 condition, they are expected to withhold responding, while in the Aware-NOGO condition, they are
332 expected to respond. Thus, catch trials will be analyzed separately for the two conditions (GO and
333 NOGO) by means of a paired sample t-test (two-tailed), in order to ensure that the behavioral
334 performance follows the above-mentioned trend. Paired sample t-tests (two-tailed) will indeed be
335 performed to test whether catch trials performance is significantly different from critical trials.

336 2.8.2 EROS data


337 Pre-processing of continuous phase delay (i.e., time-of-flight) data will be computed by means of a
338 dedicated in-house software, P-POD (Pre-Processing of Optical Data, run in Matlab MATLAB,
339 version R2013b). Thus, raw data will be normalized (i.e., corrected for phase wrapping and de-
340 trended to remove low-frequency drifts), baseline corrected and filtered by means of a 6th order
341 Butterworth band-pass filter which allows frequencies between 0.5 Hz and 15 Hz. Pulse artifact will
342 be removed by using a regression algorithm (GRATTON *et al.*, 1995). After that, data will be
343 averaged separately for each subject, condition, and channel and segmented into epochs time-locked
344 to the onset of the stimulus. Each epoch will comprise a period from 486 ms before the stimulus onset
345 to 998 ms following the stimulus onset, resulting in an epoch lasting 1484 ms. Subsequently,
346 statistical analyses will be computed with an in-house software package (Opt-3d; (Gratton, 2000)),
347 which provides statistical spatial maps of fast optical data.

348 To perform statistics, data from channels whose diffusion paths intersect a given voxel will be
349 combined (Wolf *et al.*, 2014). Phase delay data will be spatially filtered with an 8-mm Gaussian
350 kernel. Within each ROI, t-Statistics will be calculated at group level, converted into Z-scores and
351 corrected for multiple comparisons using random field theory (Worsley *et al.*, 1995; Kiebel *et al.*,
352 1999). Then, Z-scores will be weighted and orthogonally projected onto the surface of an MNI
353 template brain, according to the physical homogenous model (Arridge & Schweiger, 1995; Gratton,
354 2000).

355 In order to investigate the neural dynamics related to conscious vision and to disentangle the role of
356 the motor areas, the following contrasts between conditions will be computed: 1) Aware-GO versus
357 Unaware-GO and 2) Aware-NOGO versus Unaware-NOGO. These contrasts allow to investigate the
358 research questions the proposed study aims at answering (see Section 3 for a detailed description of
359 the planned analysis).

360 Moreover, Granger Causality analysis will be computed. Granger Causality analysis allows to explore
361 the predictive interactions between different brain areas at different time-points. Specifically, this
362 approach requires a region of interest (ROI) to be used as a “seed” and investigating whether the
363 activity of this seed predicts activity in the other ROIs at a later time-lag, by deriving statistical maps
364 from t-statistics computation (then transformed into z scores) for each lag.


365 Statistical functional analysis will be computed within specific **predetermined** regions of interest
366 (ROIs) **and time intervals**. **ROIs will be** defined by a 2-dimensional box-shaped structure, covering
367 an area of 20x20 millimeters. Critical ROIs will be selected on the basis of the results obtained in the
368 above-mentioned experiment (Colombari *et al.*, 2024) and by visual inspection of functional data. In
369 particular, they will be located in the occipital and in the left parietal and temporal lobes, specifically
370 over the primary visual cortex (V1, Brodmann Area 17), the left lateral occipital cortex (LOC,
371 Brodmann Area 19), the left supplementary motor area (SMA, Brodmann Area 6), the left premotor
372 area (PM, Brodmann Area 6) and the left primary motor cortex (M1, Brodmann Area 4). **Statistical**
373 **analysis will be computed within specific temporal windows of interest selected on the basis of the**
374 **results obtained by Colombari et al., 2024. This is to reduce the risk of false positives, as Opt3d does**
375 **not offer the possibility to correct data for multiple comparisons in the temporal domain. The specific**
376 **time windows tested for each hypothesis are listed in Table 3.**



377

378 **Figure 1. Trial procedure and stimuli:** **A)** Experimental procedure: the trial begins with a fixation cross
 379 persisting at the center of the screen for 500 ms. After that, an acoustic tone lasting 100 ms will be presented,
 380 followed by a random interval ranging from 500 to 600 ms. Then, the stimulus will be presented for 100 ms
 381 and participants will be asked to respond according to the experimental condition (i.e., Aware-GO or Aware-
 382 NOGO). **B)** Example of stimuli: on the left is shown the catch stimulus, with all the radii equally thick; on the
 383 right is depicted the critical stimulus, with the first radius, clockwise, thicker than the others.

384



385

386 **Figure 2. Covered area.** The gray area represents the area covered by the EROS montages (combined
 387 together) from the sagittal, axial and coronal point of view.

3. Study design

Question	Hypothesis	Sampling Plan	Analysis Plan	Rationale for deciding the sensitivity of the test for confirming or disconfirming the hp	Interpretation given different outcomes
<p>Q1: Can we replicate Colombari et al., 2024 findings showing that LOC is an NCC?</p>	<p>H1: We hypothesize to replicate Colombari et al., 2024 results: greater activity in LOC in an early temporal window (i.e., 150-350 ms post stimulus onset) is observed when contrasting Aware and Unaware trials in the condition in which the response is required (i.e., GO condition)</p> <p>Expected outcome: LOC aware-GO>LOC unaware-GO, as measured by EROS activity</p>	<p>Because of technical constraints of the dedicated EROS software, the effect size for EROS data is not computable and thus sample size cannot be determined basing on previous EROS findings. For this reason, the sample size estimation for the present question is basically determined according to two strategies: 1) a systematic review of existing EROS literature revealing that the typical sample size used is 12-13 participants (see Supplementary Table 1) 2) sample size estimation based on a previous EEG study of Koivisto et al., 2016, in which authors employed the same experimental paradigm adopted in the present study and aware trials were compared to unaware trials. Sample size calculation was thus performed with G-Power software (v. 3.1.9.7),</p>	<p>A1: The goal is to replicate the results of Colombari et al., 2024, in which the manual response was required for both Aware and Unaware conditions. Here, in order to perform the same analysis, early LOC activity in Aware-GO and Unaware-GO trials will be compared by using a paired-sample one-tailed t-test, computed with the EROS dedicated analysis software “Opt3d”.</p> <p>Contrast to be computed: AWARE GO VS UNAWARE GO</p> <p>ROI to be tested: LOC</p> <p>Time interval of interest: 150-350ms after stimulus onset</p>	<p>Effect size for EROS data is not computable. This is because the existing software dedicated to statistical EROS analysis (i.e., Opt-3d) does not allow to calculate this measure. However, we estimated our sample size basing of the effect size of a previous EEG study (Koivisto et al., 2016) employing the same experimental design and based on the sample used in EROS literature.</p>	<p>O1.1: A significant t-test within the interval of interest in an early time window will be interpreted as a successful replication of previous findings, supporting the involvement of LOC in NCC.</p> <p>O1.2: The absence of this effect will not confirm the hypothesis, suggesting that LOC is not involved in the conscious detection of a stimulus property.</p>

		with a power of 90% and a level of significance of 2%, resulting in 15 participants. However, since EROS signal-to-noise ratio is lower than that of EEG, we will increase our final sample to 26 24 participants, which is almost the double of the estimated sample size.			
Q2: Is the activity in LOC independent from the response?	<p>H2: We hypothesize that LOC activity is independent from response requirement: when contrasting activity elicited by Aware-NOGO trials with activity elicited by Unaware-NOGO trials, we expect to find the same activation of LOC found in the Aware-GO vs Unaware-GO contrast.</p> <p>Expected outcome:</p> <p>LOC aware-NOGO > LOC unaware-NOGO, as measured by EROS activity</p> <p>(LOC aware-GO > LOC unaware-GO) = (LOC aware-NOGO > LOC unaware-NOGO)</p>	As Q1	<p>A2.1: A paired-sample one-tailed t-test will be computed in order to compare early activity in LOC in the NOGO condition. Thus, activity in Aware-NOGO and Unaware-NOGO trials will be contrasted.</p> <p>Contrast to be computed:</p> <p>AWARE NOGO VS UNWARE NOGO</p> <p>ROI to be tested:</p> <p>LOC</p> <p>Time interval of interest:</p> <p>150-350ms after stimulus onset</p>	As above	<p>O2.1.1: A significant t-test in an an early the time window of interest will suggest that LOC activity is independent from response, since its activity is observed even when no response is required (NOGO conditions).</p> <p>O2.1.2: If greater activity in LOC in an an early the time window of interest is not observed, then it means that LOC activity is somehow related to the motor response.</p>
			A2.2: The interaction effect between awareness		O2.2.1: Significant interaction effect will suggest that activity in LOC depends from response requirement

			<p>and motor response will be tested by means of a paired-sample one-tailed t-test computed between contrast Aware-GO VS Unaware-GO and contrast Aware-NOGO VS Unaware-NOGO</p> <p>Contrast to be computed:</p> <p>(AWARE GO VS UNAWARE GO) - (AWARE NOGO VS UNAWARE NOGO)</p> <p>ROI to be tested: LOC</p> <p>Time interval of interest: 150-350ms after stimulus onset</p>		<p>O2.2.2: The absence of a difference between the two effects will suggest that motor response does not affect awareness-related activity in LOC</p>
<p>Q3: Does consciousness modulate activation of motor areas in a detection task?</p>	<p>H3: When a motor response is required, consciousness modulates activation of motor areas (MA), as activity in motor areas is triggered by LOC (Colombari et al., 2024)</p> <p>Expected outcome: MA aware-GO>MA unaware-GO, as</p>	<p>Considering that the estimated sample size for this study (n=26 24) is the double of the typical sample size of EROS studies present in literature, the same estimated sample size seems to be also adequate to answer research questions Q3 and Q4.</p>	<p>A3.1 A paired-sample one-tailed t-test will be computed in order to compare early activity in Motor Areas in the GO condition. Thus, activity in Aware-GO and Unaware-GO trials will be contrasted.</p> <p>Contrast to be computed:</p>	As above	<p>O3.1.1: A statistically significant difference between the two conditions will suggest that, even in a detection task, response related motor activity is stronger in the Aware condition compared to the Unaware one.</p> <p>In Colombari et al., 2024 this difference was observed. Importantly, in this previous study a <i>discrimination</i> task was employed and participants were asked to provide two different responses in case of Awareness (intentional) or Unawareness (random).</p>

	<p>measured by EROS activity</p> <p>LOC activity predicts MA activity (investigated by means of Granger Causality Analysis)</p>		<p>AWARE GO VS UNAWARE GO</p> <p>ROI to be tested: Motor areas</p> <p>Time interval of interest: Based on mean RTs</p>		<p>Instead, in this study participants are asked to perform a <i>detection</i> task, in which the motor behavior made to provide the response, when required, is the same for both Aware and Unaware condition and thus no response selection is required.</p> <p>O3.1.2: If no difference between the tested conditions is observed, it will suggest that in a detection task there is no difference in the motor activity related to the response.</p> <p>O3.2.1: Significant predictive interactions between LOC and motor areas will suggest that, when the stimulus enters consciousness, awareness-related activity in LOC predicts subsequent activity in motor areas. This (expected) outcome will suggest that consciousness modulates subsequent response-related motor activity, by directly triggering activation of motor areas, as observed in Colombari et al., 2024 under review</p> <p>O3.2.2: If no significant interactions between LOC and MA will be highlighted, then it would mean that activity in motor areas is not predicted by LOC. Specifically, it could be surmised that in a <i>detection</i> task, consciousness does not modulate activation of motor areas, as observed in Colombari et al., 2024 under review, where a <i>discrimination</i> task was employed. The difference in the two tasks, indeed, consists in the type of motor response required: in the case of</p>
			<p>A3.2: In order to further investigate the flow of activity occurring in the investigated brain areas, Granger Causality Analysis will be performed. In the present study, we will perform Granger analysis on the “Aware-GO VS Unaware-GO” contrast, since we are interested in investigating whether activity in motor areas is predicted by previous activity in LOC, when a motor response is required (i.e., in the GO condition). Thus, LOC will be used as seed ROI and later activity in motor areas will be investigated.</p>		

			<p>Contrast to be computed:</p> <p>AWARE GO VS UNAWARE GO</p> <p>ROI to be tested: LOC (as seed ROI) Motor areas as predicted areas</p> <p>Time interval of interest: LOC: 150-350 ms after the stimulus onset</p> <p>MA: based on mean RTs</p>		<p>the discrimination task, the participant is asked to press one button or another according to the response. Conversely, in a detection task, the participant has to press a key when the target stimulus is detected. Thus, no selection of the response is needed. This difference could play a role in the relationship between consciousness and motor areas.</p>
<p>Q4: Does consciousness modulate activation of motor areas in ABSENCE of motor response?</p>	<p>H4: Consciousness modulates activation of motor areas, even if the motor response is not required</p> <p>Expected outcome: MA aware-NOGO>MA unaware-NOGO</p> <p>LOC predicts MA (investigated by means of Granger Causality Analysis)</p>	As Q3	<p>A4.1: A paired-sample one-tailed t-test will be computed in order to compare activity in Motor Areas in the NOGO condition. Thus, activity in Aware-NOGO and Unaware-NOGO trials will be contrasted.</p> <p>Contrast to be computed:</p> <p>AWARE NOGO VS UNAWARE NOGO</p> <p>ROI to be tested: Motor areas</p> <p>Time interval of interest: Based on mean RTs</p>	As above	<p>O4.1.1: A statistically significant t-test will suggest that, when a motor response is not provided, the inhibition required to withhold responding is stronger when the visual characteristic of the stimulus is consciously perceived, compared to when no difference is perceived.</p> <p>O4.1.2: If no difference between the tested conditions is observed, this will suggest that i) no inhibition is required to withhold responding, both in the Aware and Unaware condition, or ii) the inhibition is equally strong for the two conditions.</p>

			<p>A4.2: With the aim of investigating the flow of activity occurring in the investigated brain areas also in the condition where no response is required, Granger Analysis will be performed on the “Aware-NOGO VS Unaware-NOGO” contrast. This will allow to investigate whether activity in motor areas is triggered by previous activity in LOC, even when a motor response is not required. Thus, LOC will be used as seed ROI and later activity in motor areas will be investigated.</p> <p>Contrast to be computed:</p> <p>AWARE NOGO VS UNAWARE NOGO</p> <p>ROI to be tested: LOC (as seed ROI)</p> <p>Motor areas as predicted areas</p> <p>Time interval of interest:</p>		<p>O4.2.1: If significant predictive interactions between LOC and motor areas will be observed, then consciousness modulates subsequent activity in motor areas also in absence of a motor response. This could be due to inhibition of the response processes.</p> <p>O4.2.2: If no significant interactions between LOC and MA will be highlighted, then LOC does not predict activity in motor areas in absence of motor response.</p>
--	--	--	--	--	---

			LOC: 150-350 ms after the stimulus onset MA: Based on mean RTs		
--	--	--	--	--	--

390 **4. Pilot study**

391 In order to test the experimental paradigm, we pilot-tested the task.

392 A total of 10 right-handed participants (5 females and 5 males; mean age \pm standard deviation: 21
393 years \pm 1.0) took part in the pilot study. They all reported normal or corrected-to-normal vision and
394 no history of neurological or psychiatric disorders. All of them provided written informed consent
395 before starting the experiment.

396 After the first session, two participants dropped out the experiment, hence data from 8 participants
397 were included in the statistical analyses. Moreover, in order to maintain an equal number of trials in
398 both the conditions (i.e., Aware and Unaware), the percentage of Aware and Unaware trials was
399 calculated and data from participants reporting a proportion of awareness equal or superior to 80%
400 (i.e., 3 participants) were discarded from subsequent analysis. For this pilot study, we decided to raise
401 the awareness threshold of acceptance to 80% (instead of 75%, that will be used in the experiment)
402 in order to be more inclusive, given the low number of participants.

403 Thus, in total, data from 5 participants were included in the behavioral and functional analyses.

404 **4.1 Preliminary Results**

405 *4.1.1 Behavioral results*


406 Raw data were processed by means of scripts created on Matlab (version R2017b; the MathWorks,
407 Inc., Natick, MA). According to the participants' responses, trials were sorted into the four
408 experimental conditions (i.e., Aware-GO, Unaware-NOGO, Aware-NOGO and Unaware-GO).
409 Aware trials were those trials in which the participant reported to perceive the thicker radius, while
410 Unaware trials were those trials in which participants could not perceive that the radius was thicker.
411 As specified in Section 2.8, trials with RTs lower than 150 ms or higher than 3SD from the mean
412 were removed. After removal, we had on average 830.6 trials for the Aware-GO condition, 389.2 for
413 the Unaware-NOGO condition, 738.8 trials for the Aware-NOGO condition and 491.4 for the
414 Unaware-GO.

415 Subsequently, once assessed the normality of RTs and Awareness distributions (Shapiro-Wilk test.
416 RTs distribution: $W=0.824$, $p=0.125$; Awareness distribution: $W=0.817$, $p=0.112$), the percentage of
417 Awareness for the two conditions was calculated: in the GO condition, Aware trials represented on
418 average 68.02% of the trials, while in the NOGO condition, Aware trials constituted the 59.82% of
419 the trials. Paired sample (two-tailed) t-test performed with Jamovi (version 2.3.28) highlighted that
420 there was no significant difference between the two conditions ($t_{(4)} = 1.88$, $p = .134$, Cohen's $d =$
421 $.839$), suggesting that they are comparable. Similarly, mean RTs for Aware and Unaware trials in the
422 GO condition were contrasted and the statistical analysis (Paired sample two-tailed t-test) revealed

423 that mean RTs for the Aware condition (628.530 ms) and the Unaware condition (675.317 ms) were
424 not statistically different ($t(4) = -1.77$, $p = .152$, Cohen's $d = -.791$). This indicated that there was no
425 difference in the responsiveness between the two conditions. The behavioral results are depicted in
426 Figure 3.

427 Moreover, in order to verify that the employed paradigm works as planned and that participants
428 performed the task accurately, analysis on catch trials was performed as described in Section 2.8.1
429 Behavioral data. As specified above, catch trials were those trials in which all the radii of the stimulus
430 are equally thick. Hence, in those cases, participants should report not to see the thicker radius. As
431 expected, they correctly reported not seeing the thicker radius on average the 96.47% of times
432 ($sd=2.49$) in the Aware GO condition and the 98.36% ($sd=1.89$) in the Aware NOGO condition.
433 Paired sample (two-tailed) t-test revealed no significant difference between the two conditions.

434



435

436 **Figure 3. Behavioral results.** The percentage of Awareness was calculated for both “GO” and “NOGO”
437 conditions (on the left). Mean reaction times were calculated for Aware and Unaware trials only for the “GO”
438 condition (on the right). No significant differences were observed. Error bars represent SEM and gray dots
439 represent individual data points showing the data distribution.


440 4.1.2 EROS results

441 EROS data were pre-processed with a dedicated in-house software, P-POD (Pre-Processing of Optical
442 Data, run in Matlab MATLAB, version R2013b), as described in Section 2.8. Subsequently, we
443 computed statistical analyses on pre-processed data by means of the dedicated in-house software
444 package Opt-3d.

445 For this pilot study, participants' individual structural MR images could not be acquired, so an
446 estimated MR-based head model was individually created using the Softaxic Optic system (Softaxic,
447 E.M.S., Bologna, Italy) combined with a 3D optical digitizer (Polaris Vicra, NDI, Waterloo, Canada).
448 EROS data were thus co-registered with the estimated MRI using a specific procedure performed in
449 OCP software package (as specified above). Finally, co-registered data were transformed into MNI
450 space for subsequent analyses.

451 For both GO and NOGO conditions, Aware and Unaware trials were contrasted. As shown in Figure
452 4, the Aware-GO vs Unaware-GO contrast replicated the results obtained by Colombari et al., 2024
453 ~~under review~~. In this contrast, indeed, we compared conditions in which the motor response was
454 required, thus replicating the task carried out in the previously mentioned experiment. Also in this
455 case, Aware trials elicited a sustained activation of LOC (230 and 332 ms after the stimulus onset),
456 followed by the recurrent activation of the primary visual cortex (V1) and the motor areas (MA) at
457 later stages of stimulus processing.

458 Similarly, contrasting Aware and Unaware trials in the condition where the motor response was not
459 required (i.e., the NOGO condition), greater activation of LOC was elicited in a timing comparable
460 to that of the contrast just mentioned above (i.e., 307 ms after the stimulus presentation). Interestingly,
461 also in this case awareness-related processing elicited activity in the motor areas, 563 ms after the
462 stimulus onset, despite in this condition no response was required, possibly suggesting an inhibition
463 to respond for the NOGO trials.



464 **Figure 4. EROS results.** Statistical parametric maps of the z-score difference computed EROS results
465 obtained contrasting Aware and Unaware trials in the GO (upper panel) and NOGO condition (lower panel).
466 Each map represents a 25.6 ms interval.
467

468 4.2 Preliminary Discussion

469 The aim of the present pilot study was to assess whether the task and the experimental procedure were
470 suitable to investigate the study's research questions.

471 As described in Section 4.1, the pilot study successfully replicated the trend of activations observed
472 by Colombari et al., 2024 ~~under review~~, suggesting that the proposed study proves to be feasible in
473 terms of methodology. For the sake of clarity, it is important to point out that the preliminary results
474 reported here do not reach the statistical level of significance. This outcome was expected as data

475 from only 5 participants were included in the analysis. For the same reason, we decided not to perform
476 Granger Causality analysis as for this kind of analysis results from 5 participants would have been
477 uninformative. Nevertheless, it was possible to observe that the proposed task could elicit a pattern
478 of activation similar to that observed by Colombari et al., 2024~~under review~~, suggesting that the
479 experimental paradigm proposed to investigate the research questions is suitable.

480

481 **Data availability**

482 Upon acceptance of the Stage 2 registered report, we will share all raw and processed anonymized
483 data as well as study materials publicly available as open data. Pilot raw and processed data can be
484 found on this link: https://osf.io/ebfu3/?view_only=9ec2e6bf32ba4a8bb8b858639ec40a59

485 **Code availability**

486 All analysis code will be made publicly available upon acceptance of the Stage 2 registered report.

487 **Acknowledgments**

488 The present project is supported by the grant program "Funding Consciousness Research with
489 Registered Reports" and Fondazione Cassa di Risparmio di Verona, Vicenza, Belluno e Ancona
490 "Ricerca scientifica d'eccellenza 2018", "Emergence of Consciousness: From neural dynamics to
491 complex conscious behaviour" grant no. 2018.0861; SS is supported by #NEXTGENERATIONEU
492 (NGEU) and funded by the Ministry of University and Research (MUR), National Recovery and
493 Resilience Plan (NRRP), project MNESYS (PE0000006)—"A Multiscale integrated approach to the
494 study of the nervous system in health and disease" (DN. 1553 11.10.2022), CM is supported by MIUR
495 D.M. 737/2021—"Neural correlates of perceptual awareness: from neural architecture to the
496 preservation of conscious vision in brain tumor patients".

497 **Author contributions**

498 **EC** Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data
499 Curation, Writing - Original Draft, Visualization, Funding Acquisition; **GP** Methodology, Formal
500 Analysis, Investigation, Data Curation, Writing - Review & Editing; **SM** Methodology, Investigation,
501 Writing - Review & Editing **CM** Methodology, Software, Data Curation, Writing - Review & Editing,
502 Supervision; **SS** Conceptualization, Methodology, Resources, Writing - Review & Editing,
503 Supervision, Project administration, Funding acquisition.

504 **Competing interests**

505 The authors declare no competing interests.

506 **References**

- 507 Arridge, S.R. & Schweiger, M. (1995) &title>Sensitivity to prior knowledge in optical
508 tomographic reconstruction</title> In Chance, B. & Alfano, R.R. (eds), *Optical*
509 *Tomography, Photon Migration, and Spectroscopy of Tissue and Model Media: Theory,*
510 *Human Studies, and Instrumentation*. SPIE, pp. 378–388.
- 511 Aru, J., Bachmann, T., Singer, W., & Melloni, L. (2012) Distilling the neural correlates of
512 consciousness. *Neurosci. Biobehav. Rev.*, **36**, 737–746.
- 513 Baniqued, Pauline L.; Low, Kathy A.; Fabiani, Monica; Gratton, G. (2013) Frontoparietal Traffic
514 Signals: A Fast Optical Imaging Study of Preparatory Dynamics in Response Mode Switching.
515 *J. Cogn. Neurosci.* 25 887–902.,.
- 516 Bola, M. & Doradzińska, Ł. (2021) Perceptual Awareness Negativity—Does It Reflect Awareness
517 or Attention? *Front. Hum. Neurosci.*, **15**, 1–4.
- 518 Chiarelli, A.M., Maclin, E.L., Low, K.A., Fabiani, M., & Gratton, G. (2015) Comparison of
519 procedures for co-registering scalp-recording locations to anatomical magnetic resonance
520 images. *J. Biomed. Opt.*, **20**, 016009.
- 521 Ciupińska, K., Orłowska, W., Zębrowski, A., Łępa, L., Koculak, M., Bola, M., & Wierzchon, M.
522 (2024) The influence of spatial and temporal attention on visual awareness—a behavioral and
523 ERP study. *Cereb. Cortex*, **34**.
- 524 Cohen, M.A., Ortego, K., Kyroudis, A., & Pitts, M. (2020) Distinguishing the Neural Correlates of
525 Perceptual Awareness and Postperceptual Processing. *J. Neurosci.*, **40**, 4925–4935.
- 526 Colombari, E., Parisi, G., Tafuro, A., Mele, S., Mazzi, C., & Savazzi, S. (2024) Beyond primary
527 visual cortex: The leading role of lateral occipital complex in early conscious visual
528 processing. *Neuroimage*, **298**, 120805.
- 529 Dellert, T., Krebs, S., Bruchmann, M., Schindler, S., Peters, A., & Straube, T. (2022) Neural
530 correlates of consciousness in an attentional blink paradigm with uncertain target relevance.
531 *Neuroimage*, **264**, 119679.
- 532 Dellert, T., Müller-Bardorff, M., Schlossmacher, I., Pitts, M., Hofmann, D., Bruchmann, M., &
533 Straube, T. (2021) Dissociating the neural correlates of consciousness and task relevance in
534 face perception using simultaneous EEG-fMRI. *J. Neurosci.*, **41**, 7864–7875.

- 535 Dembski, C., Koch, C., & Pitts, M. (2021) Perceptual awareness negativity: a physiological
536 correlate of sensory consciousness. *Trends Cogn. Sci.*, **25**, 660–670.
- 537 Doradzińska, Ł. & Bola, M. (2024) Early Electrophysiological Correlates of Perceptual
538 Consciousness Are Affected by Both Exogenous and Endogenous Attention. *J. Cogn.
539 Neurosci.*, **36**, 1297–1324.
- 540 Förster, J., Koivisto, M., & Revonsuo, A. (2020) ERP and MEG correlates of visual consciousness:
541 The second decade. *Conscious. Cogn.*, **80**.
- 542 Gratton, G. (2000) “Opt-cont” and “opt-3D”: A software suite for the analysis and 3D
543 reconstruction of the event-related optical signal (EROS). *Psychophysiology*,.
- 544 Gratton, G. & Corballis, P.M. (1995) Removing the heart from the brain: Compensation for the
545 pulse artifact in the photon migration signal. *Psychophysiology*, **32**, 292–299.
- 546 GRATTON, G., CORBALLIS, P.M., CHO, E., FABIANI, M., & HOOD, D.C. (1995) Shades of
547 gray matter: Noninvasive optical images of human brain reponses during visual stimulation.
548 *Psychophysiology*,.
- 549 Gratton, G. & Fabiani, M. (1998) Dynamic brain imaging: Event-related optical signal (EROS)
550 measures of the time course and localization of cognitive-related activity. *Psychon. Bull. Rev.*,
551 **5**, 535–563.
- 552 Gratton, G. & Fabiani, M. (2001) Shedding light on brain function: The event-related optical signal.
553 *Trends Cogn. Sci.*, **5**, 357–363.
- 554 Gratton, G. & Fabiani, M. (2003) The event-related optical signal (EROS) in visual cortex:
555 Replicability, consistency, localization, and resolution. *Psychophysiology*, **40**, 561–571.
- 556 Gratton, G. & Fabiani, M. (2010) Fast optical imaging of human brain function. *Front. Hum.
557 Neurosci.*, **4**, 1–9.
- 558 Gratton, G., Fabiani, M., Corballis, P.M., Hood, D.C., Goodman-Wood, M.R., Hirsch, J., Kim, K.,
559 Friedman, D., & Gratton, E. (1997) Fast and localized event-related optical signals(EROS) in
560 the human occipital cortex: Comparisons with the visual evoked potential and fMRI.
561 *Neuroimage*, **6**, 168–180.
- 562 Gratton, G., Goodman-Wood, M.R., & Fabiani, M. (2001) Comparison of neuronal and
563 hemodynamic measures of the brain response to visual stimulation: An optical imaging study.
564 *Hum. Brain Mapp.*, **13**, 13–25.

565 Gratton, G., Low, K.A., Maclin, E.L., Brumback, C.R., Gordon, B., & Fabiani, M. (2006) Time
566 course of activation of human occipital cortex measured with the event-related optical signal
567 (EROS). *Opt. InfoBase Conf. Pap.*, 5–8.

568 Gratton, G., Sarno, A., Maclin, E., Corballis, P.M., & Fabiani, M. (2000) Toward noninvasive 3-D
569 imaging of the time course of cortical activity: Investigation of the depth of the event-related
570 optical signal. *Neuroimage*, **11**, 491–504.

571 Hatamimajoumerd, E., Ratan Murty, N.A., Pitts, M., & Cohen, M.A. (2022) Decoding perceptual
572 awareness across the brain with a no-report fMRI masking paradigm. *Curr. Biol.*, **32**, 4139-
573 4149.e4.

574 Hense, A., Peters, A., Bruchmann, M., Dellert, T., & Straube, T. (2024) Electrophysiological
575 correlates of sustained conscious perception. *Sci. Rep.*, **14**, 1–11.

576 Kiebel, S.J., Poline, J.B., Friston, K.J., Holmes, A.P., & Worsley, K.J. (1999) Robust smoothness
577 estimation in statistical parametric maps using standardized residuals from the general linear
578 model. *Neuroimage*, **10**, 756–766.

579 Knight, R.S., Chen, T., Gratton, G., Fabiani, M., Savazzi, S., Mazzi, C., & Beck, D.M. (2024)
580 Neuropsychologia Bypassing input to V1 in visual awareness : A TMS-EROS investigation.
581 *Neuropsychologia*, **198**, 108864.

582 Koivisto, M., Lähteenmäki, M., Sørensen, T.A., Vangkilde, S., Overgaard, M., & Revonsuo, A.
583 (2008) The earliest electrophysiological correlate of visual awareness? *Brain Cogn.*, **66**, 91–
584 103.

585 Koivisto, M. & Revonsuo, A. (2010) Event-related brain potential correlates of visual awareness.
586 *Neurosci. Biobehav. Rev.*, **34**, 922–934.

587 Koivisto, M., Revonsuo, A., & Lehtonen, M. (2006) Independence of visual awareness from the
588 scope of attention: An electrophysiological study. *Cereb. Cortex*, **16**, 415–424.

589 Koivisto, M., Salminen-Vaparanta, N., Grassini, S., & Revonsuo, A. (2016) Subjective visual
590 awareness emerges prior to P3. *Eur. J. Neurosci.*, **43**, 1601–1611.

591 Kronemer, S.I., Aksen, M., Ding, J.Z., Ryu, J.H., Xin, Q., Ding, Z., Prince, J.S., Kwon, H., Khalaf,
592 A., Forman, S., Jin, D.S., Wang, K., Chen, K., Hu, C., Agarwal, A., Saberski, E., Mohammad,
593 S., Wafa, A., Morgan, O.P., Wu, J., Christison-lagay, K.L., Hasulak, N., Morrell, M., Urban,
594 A., Constable, R.T., & Pitts, M. (2022) Human visual consciousness involves large scale

595 cortical and subcortical networks independent of task report and eye movement activity.

596 Leung, P., Franconeri, S., Grabowecky, M., & Suzuki, S. (2009) Rapid eye-fixation training
597 without eyetracking **16**, 491–496.

598 Liu, Y., Paradis, A.L., Yahia-Cherif, L., & Tallon-Baudry, C. (2012) Activity in the lateral occipital
599 cortex between 200 and 300 ms distinguishes between physically identical seen and unseen
600 stimuli. *Front. Hum. Neurosci.*, **6**, 1–9.

601 Low, K.A., Leaver, E., Kramer, A.F., Fabiani, M., & Gratton, G. (2006) Fast optical imaging of
602 frontal cortex during active and passive oddball tasks. *Psychophysiology*, **43**, 127–136.

603 Makeig, S. & Jung, T. (2000) Independent Component Analysis of Simulated ERP Data. *Brain*, 1–
604 24.

605 Mazzi, C., Mazzeo, G., & Savazzi, S. (2020) Late Positivity Does Not Meet the Criteria to be
606 Considered a Proper Neural Correlate of Perceptual Awareness. *Front. Syst. Neurosci.*, **14**, 1–
607 14.

608 Medvedev, A. V., Kainerstorfer, J., Borisov, S. V., Barbour, R.L., & VanMeter, J. (2008) Event-
609 related fast optical signal in a rapid object recognition task: Improving detection by the
610 independent component analysis. *Brain Res.*, **1236**, 145–158.

611 Medvedev, A. V., Kainerstorfer, J.M., Borisov, S. V., Gandjbakhche, A.H., & VanMeter, J. (2010)
612 “Seeing” electroencephalogram through the skull: imaging prefrontal cortex with fast optical
613 signal. *J. Biomed. Opt.*, **15**, 061702.

614 Oldfield, R. (1971) The assessment and analysis of handedness: the Edinburgh Inventory.
615 *Neuropsychologia*,.

616 Parisi, G., Mazzi, C., Colombari, E., Chiarelli, A.M., Metzger, B.A., Marzi, C.A., & Savazzi, S.
617 (2020) Spatiotemporal dynamics of attentional orienting and reorienting revealed by fast
618 optical imaging in occipital and parietal cortices. *Neuroimage*, **222**, 117244.

619 Pitts, M.A., Padwal, J., Fennelly, D., Martínez, A., & Hillyard, S.A. (2014) Gamma band activity
620 and the P3 reflect post-perceptual processes, not visual awareness. *Neuroimage*, **101**, 337–350.

621 Proulx, N., Samadani, A.A., & Chau, T. (2018) Quantifying fast optical signal and event-related
622 potential relationships during a visual oddball task. *Neuroimage*, **178**, 119–128.

623 Railo, H., Revonsuo, A., & Koivisto, M. (2015) Behavioral and electrophysiological evidence for
624 fast emergence of visual consciousness. *Neurosci. Conscious.*, **2015**, 1–12.

- 625 Schelonka, K., Grauly, C., Canseco-gonzalez, E., & Pitts, M.A. (2017) ERP signatures of
626 conscious and unconscious word and letter perception in an inattentional blindness paradigm.
627 *Conscious. Cogn.*, **54**, 56–71.
- 628 Schlossmacher, I., Dellert, T., Pitts, M., Bruchmann, M., & Straube, T. (2020) Differential effects
629 of awareness and task relevance on early and late ERPs in a no-report visual oddball paradigm.
630 *J. Neurosci.*, **40**, 2906–2913.
- 631 Seth, A.K., Barrett, A.B., & Barnett, L. (2015) Granger causality analysis in neuroscience and
632 neuroimaging. *J. Neurosci.*, **35**, 3293–3297.
- 633 Shafto, J.P. & Pitts, M.A. (2015) Neural signatures of conscious face perception in an inattentional
634 blindness paradigm. *J. Neurosci.*, **35**, 10940–10948.
- 635 Toscano, J.C., Anderson, N.D., Fabiani, M., Gratton, G., & Garnsey, S.M. (2018) The time-course
636 of cortical responses to speech revealed by fast optical imaging. *Brain Lang.*, **184**, 32–42.
- 637 Tse, C.Y. & Penney, T.B. (2007) Preattentive change detection using the event-related optical
638 signal. *IEEE Eng. Med. Biol. Mag.*, **26**, 52–58.
- 639 Tse, C.Y., Shum, Y.H., Xiao, X.Z., & Wang, Y. (2021) Fronto-occipital mismatch responses in pre-
640 attentive detection of visual changes: Implication on a generic brain network underlying
641 Mismatch Negativity (MMN). *Neuroimage*, **244**, 118633.
- 642 Tsuchiya, N., Wilke, M., Frässle, S., & Lamme, V.A.F. (2015) No-Report Paradigms: Extracting
643 the True Neural Correlates of Consciousness. *Trends Cogn. Sci.*, **19**, 757–770.
- 644 Vanni, S., Revonsuo, A., Saarinen, J., & Hari, R. (1996) Visual awareness of objects correlates with
645 activity of right occipital cortex. *Neuroreport*, **8**, 183–186.
- 646 Wolf, M., Wolf, U., Choi, J.H., Toronov, V., Paunescu, L.A., Michalos, A., & Gratton, E. (2003)
647 Fast cerebral functional signal in the 100-ms range detected in the visual cortex by frequency-
648 domain near-infrared spectrophotometry. *Psychophysiology*, **40**, 521–528.
- 649 Wolf, U., Wolf, M., Toronov, V., Michalos, A., Paunescu, L.A., & Gratton, E. (2014) Detecting
650 cerebral functional slow and fast signals by frequency-domain near-infrared spectroscopy
651 using two different sensors. In *Biomedical Optical Spectroscopy and Diagnostics (2000)*,
652 *Paper TuF10*. The Optical Society, p. TuF10.
- 653 Worsley, K.J., Poline, J.B., Vandal, A.C., & Friston, K.J. (1995) Tests for distributed, nonfocal
654 brain activations. *Neuroimage*, **2**, 183–194.

655 Ye, M., Wang, A., Liang, H., & Liu, X. (2024) Late positivity correlates with subjective reports:
656 Evidence from the low-frequency and high-frequency reporting tasks. *Neuroscience*,.

657